

Supplementary Table S1. Pearson correlation coefficients between TSE (mg/infusion), SG (mg/infusion), PC (mg GAE/infusion), TC (mg GAE/infusion), FC (mg QE/infusion), and IC₅₀ (µg/mL)

Response	RbA	SG	TSE	PC	TC	FC	IC ₅₀
RbA	1.000000	0.998778	0.774173	0.823399	0.491993	0.819738	-0.466552
	(*)	(0.000000) ^a	(0.000000)	(0.000000)	0.002306)	(0.000000)	0.004127)
SG	0.998778	1.000000	0.775130	0.835278	0.507140	0.817801	-0.459380
	(0.000000) ^a	(*)	(0.000000)	(0.000000)	(0.001596)	(0.000000)	(0.004826)
TSE	0.774173	0.775130	1.000000	0.757830	0.532014	0.778275	-0.364069
	(0.000000)	(0.000000)	(*)	(0.000000)	(0.000839)	(0.000000)	(0.029047)
PC	0.823399	0.835278	0.757830	1.000000	0.751657	0.828876	-0.608258
	(0.000000)	(0.000000)	(0.000000)	(*)	(0.000000)	(0.000000)	(0.000083)
TC	0.491993	0.507140	0.532014	0.751657	1.000000	0.603060	-0.298665
	(0.002306)	(0.001596)	(0.000839)	(0.000000)	(*)	(0.000099)	(0.076831)
FC	0.819738	0.817801	0.778275	0.828876	0.603060	1.000000	-0.694845
	(0.000000)	(0.000000)	(0.000000)	(0.000000)	(0.000099)	(*)	(0.000003)
IC₅₀	-0.466552	-0.459380	-0.364069	-0.608258	-0.298665	-0.694845	1.000000
	(0.004127)	(0.004826)	(0.029047)	(0.000083)	(0.076831)	(0.000003)	(*)

^a P-values are indicates in parentheses (95% CL, based on 36 assays with 2 replicates each)