

Table S2 The pH value of BBJ samples during storage.

Samples	4°C				25°C			
	0 d	30 d	60 d	90 d	0 d	30 d	60 d	90 d
Control	3.68 ±	3.54 ±	3.45 ±	3.28 ±	3.68 ±	3.40 ±	3.57 ±	3.67 ±
	0.03 b	0.01 b	0.02 b	0.00 c	0.03 b	0.01 f	0.01 b	0.01 a
Ste2	3.65 ±	3.56 ±	3.39 ±	3.42 ±	3.65 ±	3.45 ±	3.24 ±	3.65 ±
	0.03 b	0.01 b	0.01 d	0.01 a	0.03 b	0.01 d	0.02 e	0.01 b
Ste4	3.62 ±	3.50 ±	3.38 ±	3.30 ±	3.62 ±	3.47 ±	3.54 ±	3.65 ±
	0.02 bc	0.01 b	0.01 d	0.03 b	0.02 bc	0.01c	0.01 b	0.00 b
Ste8	3.60 ±	3.57 ±	3.38 ±	3.18 ±	3.60 ±	3.47 ±	3.56 ±	3.64 ±
	0.03 c	0.02 b	0.02 d	0.02 g	0.03 c	0.01c	0.03 b	0.01 b
Ste12	3.57 ±	3.50 ±	3.35 ±	3.13 ±	3.57 ±	3.62 ±	3.56 ±	2.91 ±
	0.03 cd	0.01 b	0.01 g	0.01 h	0.03 cd	0.02 b	0.02 b	0.01 g
Ste16	3.61 ±	3.49 ±	3.37 ±	3.19 ±	3.61 ±	3.39 ±	3.54 ±	3.52 ±
	0.02 c	0.00 b	0.01 e	0.00 f	0.02 c	0.01 f	0.01bc	0.01 d
Ste20	3.58 ±	3.47 ±	3.44 ±	3.25 ±	3.58 ±	3.86 ±	3.50 ±	2.95 ±
	0.00 cd	0.01 b	0.02 c	0.02 d	0.00 cd	0.01 a	0.02 d	0.01 f
Sucralose	3.67 ±	3.52 ±	3.46 ±	3.21 ±	3.67 ±	3.43 ±	3.64 ±	3.31 ±
	0.01 b	0.02 a	0.01 a	0.01 e	0.01 b	0.01 e	0.01 a	0.01 e
Sucrose	3.72 ±	3.49 ±	3.36 ±	3.00 ±	3.72 ±	3.38 ±	2.95 ±	3.59 ±
	0.03 a	0.03 b	0.01 f	0.01 i	0.03 a	0.01 fg	0.02 f	0.01 c

Different letters in the upper-corner indicated significant difference among different groups in the same time period and storage condition ($P < 0.05$).