TABLE S2 | Pairwise mitochondrial F_{ST} values and least-cost path distances among the 16 sampling localities (populations) of *P. przewalskii*. The upper triangle presents least-cost path distances, whereas the lower triangle displays pairwise F_{ST} values. Locality abbreviations are consistent with those provided in Figure 1.

	WQ	DQ	HQ	YL	QQ	JL	ZQ	DZ	YQ	ZW	JT	WW	SQ	SW	LB	LZ
WQ	-	71.60	43.93	150.79	102.78	104.10	110.14	163.72	214.50	125.54	139.17	165.20	258.84	99.62	229.11	238.38
DQ	0.33	-	64.70	139.31	147.69	151.05	157.08	210.66	259.45	170.49	184.12	210.15	270.43	156.39	274.05	283.32
HQ	0.25	0.26	-	111.11	86.33	98.23	103.10	156.68	204.64	115.68	129.31	155.34	299.72	114.39	219.24	228.51
YL	0.71	0.69	0.79	-	108.73	169.80	160.35	209.33	256.43	166.16	179.85	207.13	401.02	216.33	271.04	280.31
QQ	0.08	0.48	0.48	0.75	-	81.71	59.49	102.27	148.69	58.07	71.76	99.39	361.16	153.17	163.29	172.57
JL	0.44	0.47	0.49	0.70	0.50	-	36.81	87.80	150.72	64.40	75.71	101.12	362.48	133.81	165.03	174.30
ZQ	0.32	0.43	0.46	0.59	0.29	0.39	-	58.88	116.71	30.67	42.00	67.41	368.52	150.29	131.31	140.58
DZ	0.88	0.91	0.93	0.97	0.89	0.82	0.49	-	85.85	53.50	51.44	50.43	422.10	203.87	102.05	111.32
YQ	0.86	0.89	0.91	0.96	0.87	0.80	0.47	0.55	-	92.95	81.50	49.86	472.88	263.95	36.93	43.39
ZW	0.86	0.88	0.90	0.94	0.86	0.79	0.48	0.42	0.51	-	14.37	43.65	383.92	175.56	107.55	116.83
JT	0.83	0.86	0.88	0.93	0.84	0.77	0.43	0.34	0.34	0.12	-	32.20	397.55	189.19	96.10	105.37
WW	0.84	0.87	0.89	0.93	0.85	0.78	0.47	0.44	0.14	0.40	0.24	-	423.58	214.64	64.46	73.73
SQ	0.91	0.94	0.96	0.99	0.92	0.87	0.67	0.94	0.91	0.89	0.86	0.85	-	336.37	487.48	496.76
SW	0.91	0.93	0.95	0.99	0.91	0.85	0.66	0.94	0.90	0.89	0.86	0.85	0.97	-	278.55	287.82
LB	0.94	0.95	0.96	0.98	0.94	0.91	0.83	0.97	0.96	0.95	0.94	0.95	0.98	0.98	-	16.82
LZ	0.58	0.59	0.62	0.66	0.58	0.53	0.34	0.43	0.36	0.41	0.36	0.39	0.55	0.53	0.52	-