

Table S4 The mobbing times and individual number of the top ten mobbing bird species at different elevations.

1800-2200 m elevational band					2800-3200 m elevational band				
Species	Mobbing number	MF _{low} (%)	Mobbing number of individuals	Ratio (%)	Species	Mobbing number	MF _{high} (%)	Mobbing number of individuals	Ratio (%)
<i>Parus monticolus</i>	25	56.82	54	17.48	<i>Phylloscopusforresti</i>	26	53.06	79	13.46
<i>Sitta nagaensis</i>	14	31.82	23	7.44	<i>Lophophanesdichrous</i>	23	46.94	52	8.86
<i>Aethopyga gouldiae</i>	11	25.00	26	8.41	<i>Sylviparus modestus</i>	22	44.90	42	7.16
<i>Periparusvenustus</i>	10	22.73	18	5.83	<i>Phylloscopusclaudiae</i>	20	40.82	57	9.71
<i>Culicicapaceylonensis</i>	8	18.18	16	5.18	<i>Periparusater</i>	19	38.78	62	10.56
<i>Phylloscopusclaudiae</i>	7	15.91	14	4.53	<i>Ficedula strophinata</i>	18	36.73	31	5.28
<i>Phylloscopustephrocephalus</i>	6	13.64	11	3.56	<i>Parus monticolus</i>	15	30.61	28	4.77
<i>Pomatorhinus ruficollis</i>	6	13.64	10	3.24	<i>Yuhina gularis</i>	12	24.49	25	4.26
<i>Dicaeum ignipectus</i>	6	13.64	8	2.59	<i>Pericrocotusethologus</i>	12	24.49	23	3.92
<i>Yuhina gularis</i>	5	11.36	15	4.85	<i>Phylloscopusvalentini</i>	12	24.49	19	3.24

Note: Boldfaced species are those shared across different elevations.