

Table 2S. Drying dynamics of wet mycelial biomass (g) of *Fomes inzengae* strains at room temperature to constant weight during four days (D₁–D₄).

Strain	Host tree	D1	D2	D3	D4
Fi/3	<i>C. betulus</i>	8.55 ± 0.05	3.35 ± 0.15	0.85 ± 0.05	0.4 ± 0.02
Fi/26	<i>C. betulus</i>	8.05 ± 0.45	3.95 ± 0.05	2.15 ± 0.05	0.4 ± 0.02
Fi/24	<i>F. orientalis</i>	8.05 ± 0.55	2.4 ± 0.22	0.71 ± 0.11	0.4 ± 0.01
Fi/30	<i>F. orientalis</i>	8.3 ± 0.5	3.65 ± 0.05	1.02 ± 0.13	0.4 ± 0.01
Fi/15	<i>S. alba</i>	8.8 ± 0.7	3.3 ± 0.2	1.35 ± 0.55	0.45 ± 0.05
Fi/16	<i>S. alba</i>	8.3 ± 0.67	2.9 ± 0.42	1.25 ± 0.15	0.4 ± 0.02
Fi/8	<i>J. regia</i>	7.65 ± 0.25	3.1 ± 0.61	0.95 ± 0.05	0.4 ± 0.02
Fi/20	<i>P. nigra</i>	9.95 ± 0.35	3.5 ± 0.22	0.83 ± 0.23	0.55 ± 0.05
Fi/25	<i>P. nigra</i>	9.85 ± 0.25	4.4 ± 0.51	1.61 ± 0.31	0.55 ± 0.05
Fi/31	<i>A. platanoides</i>	9.55 ± 0.45	5.2 ± 0.63	2.52 ± 0.12	0.55 ± 0.05