

Supplementary Table S2 The primer sequences used for qPCR assay

Gene	Forward primer (5' to 3')	Reverse primer (3' to 5')
<i>Acta2</i>	GGACGTACAACCTGGTATTGTGC	TCGGCAGTAGTCACGAAGGA
<i>ACTA2</i>	CTATGAGGGCTATGCCTTGCC	GCTCAGCAGTAGTAACGAAGGA
<i>Actb</i>	GACGTTGACATCCGTAAAGACC	AGCCAGAGCAGTAATCTCC
<i>Bsep</i>	TCTGACTCAGTGATTCTTCGCA	CCCATAAACATCAGCCAGTTGT
<i>Cd1d</i>	GGCCTGATCGTCTTCATAGTA	GTCACCGGATGTCTTGATAAGC
<i>Colla1</i>	CTGGCGGTTCAAGTCCAAT	TTCCAGGCAATCCACGAGC
<i>COL1A1</i>	GAGGGCCAAGACGAAGACATC	CAGATCACGTCATCGCACAAAC
<i>Colla2</i>	GGTGAGCCTGGTCAAACGG	ACTGTGTCCTTTCACGCCTTT
<i>Ctss</i>	AGTGGGCATGAACGATATGGG	GTCAGGCAATGTCCGATTAGAG
<i>CTSS</i>	TGACAACGGCTTTCAGTACA	GGCAGCACGATATTTTGAGTCAT
<i>Cyp7a1</i>	AAACTCCCTGTCATACCACAAAG	TTCCATCACTTGGGTCTATGC
<i>Cyp27a1</i>	AGGGCAAGTACCCAATAAGAGA	TCGTTTAAGGCATCCGTGTAGA
<i>Cyp2b10</i>	GACTTTGGGATGGGAAAGAG	CCAAACACAATGGAGCAGAT
<i>Fxr</i>	AGGTCGGTGTGAACGGATTTG	GGGGTCGTTGATGGCAACA
<i>Il12</i>	CTCCCTTGGATCTGAGCTGGA	GTTGGAACGCTGACCATAGAGA
<i>Mrp3</i>	GACTCTCATGTGGCGAAGCA	TACCAGCACCCGAGTCTTGC
<i>Ntcp</i>	ACCCTACGTCCTCAAGGCAG	AGCCAGTAAGTGTGGTGTCA
<i>Oatp</i>	CAATTCGGTATCCCCACGTCT	CTGCACATCCTACACCAATCAT
<i>Tgfb1</i>	CTTCAATACGTCAGACATTTCGGG	GTAACGCCAGGAATTGTTGCTA