

Supplementary Material II. Aggregated (Agg.), regular (Reg.), or random (Ran.) distribution (Dist.) of the solution sources on 48 *Sapindus saponaria* (Sapindaceae) saplings

Solution Sources	Chi-square test			
	Var.*	Mean	P	Dist.
Ara.: Araneidae	1.97	0.72	0.00	Agg.
Anyphaenidae, <i>Teudis</i> sp.	0.07	0.07	0.56	Ran.
Oxyopidae	0.32	0.33	0.50	Ran.
<i>Oxyopes salticus</i> Hentz, 1845	0.02	0.02	0.47	Ran.
Salticidae	0.55	0.30	0.00	Agg.
<i>Aphirape uncifera</i> Tullgren, 1905	0.07	0.07	0.56	Ran.
<i>Uspachus</i> sp.	0.09	0.09	0.60	Ran.
Sparassidae, <i>Quemedice</i> sp.	0.07	0.07	0.56	Ran.
Tetragnathidae, <i>Leucauge</i> sp.	2.36	0.30	0.00	Agg.
Thomisidae, <i>Tmarus</i> sp.	0.05	0.05	0.51	Ran.
<i>Aphantochilus rogersi</i> O. Pic.-Cam., 1871	0.09	0.09	0.60	Ran.
Col.: Coccinellidae, <i>Cycloneda sanguinea</i> L., 1763	0.11	0.07	0.01	Agg.
Dip.: Dolichopodidae	0.36	0.21	0.00	Agg.
Syrphidae, <i>Syrphus</i> sp.	0.05	0.05	0.51	Ran.
Hym.: Apidae, <i>Tetragonisca angustula</i> Latreille, 1811	0.09	0.05	0.00	Agg.
Formicidae, <i>Brachymyrmex</i> sp.	55.30	4.28	0.00	Agg.
<i>Camponotus</i> sp.	39.07	3.02	0.00	Agg.
<i>Ectatoma</i> sp.	1.06	0.47	0.00	Agg.
<i>Pheidole</i> sp.	337.08	6.33	0.00	Agg.
<i>Pseudomyrmex termitarius</i> (Smith, 1877)	32.25	2.81	0.00	Agg.
Vespidae, <i>Polybia</i> sp.	0.27	0.14	0.00	Agg.
Mant.: Mantidae, <i>Mantis religiosa</i> L., 1758	0.29	0.26	0.25	Ran.
Neur.: Chrysopidae, <i>Chrysoperla</i> sp.	0.11	0.07	0.01	Agg.

*= variance.